Capillary Blood Lead Testing Form

NAPA-SOLANO-YOLO-MARIN-MENDOCINO COUNTY PUBLIC HEALTH LABORATORY

2201 COURAGE DRIVE MS 9-200, FAIRFIELD, CA 94533; TELEPHONE (707) 784-4410; FAX (707) 423-1979 www.solanocounty.com/depts/ph/bureaus/laboratory

SUBMITTER INFORMATION				The state of the s
Organization name:				OUNT Votended 1850
Address:				
Phone:		Fax:		COUVEY OF MARIN V
PATIENT DEMOGRAPHIC	CS—PLEASE FILL CO	MPLETELY (Refer to web	site for race and ethr	nicity responses)
Last name	First name		Birthdate	Date collected
Address (#, street, city, zip)		Pregnancy status ☐ Pregnant ☐ Not pregnant ☐ Unknown	□ Male □ Female □	Time collected
Telephone no.		Race	Ethnicity	
Employer name (if applicable)		Employer telephone	Employer address	
TEST SITE INFORMATIO	N— PLEASE FILL CO	MPLETELY AND CLEAR	LY	
Practitioner name, NPI #	‡		Accession #	ICD-10-CM Code:
BILLING INFORMATION-	-PLEASE CHECK BOX	X FOR BILLING SOURC	E AND WRITE-IN A	CCOUNT NUMBER
ATTACH COPY OF PATIE	ENT IDENTIFICATION	CARD FOR TYPE OF CO	VERAGE	
☐ Submitter	☐ Partnersh	nip (PHC) NUM	IBER:	
☐ CHDP	☐ Other:	☐ Other:		
☐ Medi-Cal				
SPECIMEN TYPE/SOURC	E—PLEASE CHECK	APPROPRIATE BOX		
☐ Blood fingerstick (cap	Drawer's in	itials:		
TEST INFORMATION				
COMMENTS/SPECIAL IN	STRUCTIONS:			

SPECIMEN REQUIREMENTS

<u>FINGERSTICK (CAPILLARY)</u>: Suggest using an adult lancet for finger puncture. Touch the capillary tube to the blood drop, and fill to the 50-microliter black line. Inspect the tube for proper filling. Place the capillary tube into the Treatment Reagent Vial, then insert the plunger into the top of the capillary tube and push down, ensuring that the entire volume of blood is dispensed. Replace the vial cap. Invert the vial 8 to 10 times to mix completely. Store refrigerated. The mixture of blood and treatment reagent is stable for up to 7 days if refrigerated.

DATE/TIME RECEIVED